

Package Leaflet: Information for the patient

EPANUTIN® 30 mg/ 5ml Oral Suspension (phenytoin)

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Epanutin is and what it is used for
2. What you need to know before you take Epanutin
3. How to take Epanutin
4. Possible side effects
5. How to store Epanutin
6. Contents of the pack and other information

1. What Epanutin is and what it is used for

This medicine contains phenytoin, which is one of a group of medicines called anti-epileptic drugs; these medicines are used to treat epilepsy.

Epanutin can be used to control epilepsy, to control or prevent seizures during or after brain surgery or severe head injury. Epanutin can also be used to treat trigeminal neuralgia (facial nerve pain).

You should consult your doctor if you are unsure why you have been given Epanutin 30mg/5ml Oral Suspension if you do not feel better or if you feel worse.

2. What you need to know before you take Epanutin

Do not take Epanutin

- if you are allergic (hypersensitive) to phenytoin, or any of the other ingredients of this medicine (listed in section 6)
- if you are allergic to other medicines for epilepsy
- if you are also taking delavirdine (used for HIV therapy).

Warnings and precautions

Talk to your doctor or pharmacist before you take Epanutin if you suffer from or have suffered in the past from any of the following conditions:

- Liver disease
- Kidney disease
- Porphyrria (an inherited disease that affects haemoglobin biosynthesis)
- Alcohol dependence.

Epanutin can cause problems with your heart, including a slow heartbeat. Let your healthcare provider know right away if you have these symptoms.

You should be administered Epanutin with caution if you suffer from kidney or liver problems.

A small number of people being treated with antiepileptics such as phenytoin have had thoughts of harming or killing themselves. If at any time you have these thoughts, immediately contact your doctor.

There is a risk of harm to the unborn child if Epanutin is used during pregnancy. Women of childbearing age should use effective contraception with Epanutin (see Pregnancy, contraception in women, and breast-feeding).

Potentially life-threatening skin rashes (Stevens Johnson syndrome, toxic epidermal necrolysis) have been reported with the use of Epanutin, appearing initially as reddish target-like spots or circular patches often with central blisters on the trunk. Additional signs to look for include ulcers in the mouth, throat, nose, genitals and conjunctivitis (red and swollen eyes). These potentially life-threatening skin rashes are often accompanied by flu-like symptoms. The rash may progress to widespread blistering or peeling of the skin. The highest risk for occurrence of serious skin reactions is within the first weeks of treatment. If you have developed Stevens-Johnson syndrome or toxic epidermal necrolysis with the use of Epanutin, you must not be re-started on Epanutin at any time.

If you develop a rash or these skin symptoms, stop taking Epanutin, seek urgent advice from a doctor and tell him that you are taking this medicine. Consult your doctor before discontinuing Epanutin. If you suddenly stop taking this medicine you may have a seizure.

This risk of these serious skin side effects may be associated with a variant in genes in a subject with Chinese or Thai origin. If you are of such origin and have been tested previously carrying this genetic variant (HLA-B*1502), discuss this with your doctor before taking Epanutin.

Black patients may be at greater risk of liver problems, serious skin reactions and allergic reactions.

Other medicines and Epanutin

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Some medicines can affect the way Epanutin works, or Epanutin itself can reduce the effectiveness of other medicines taken at the same time. These include (Not all medicines are listed here. Talk with your doctor or pharmacist):

- Medicines used for heart and circulation problems (e.g. dicoumarol, digitoxin, digoxin, disopyramide, mexiletine, nisoldipine, amiodarone, furosemide, quinidine, reserpine, warfarin, and calcium channel blockers including diltiazem and nifedipine)
- Medicines used for epilepsy (e.g. carbamazepine, lamotrigine, phenobarbital, sodium valproate and valproic acid, topiramate, oxcarbazepine, succinimides including ethosuximide and vigabatrin)
- Medicines used to treat fungal infections (e.g. amphotericin B, fluconazole, itraconazole, ketoconazole, posaconazole, voriconazole and miconazole)

- Medicines used for tuberculosis and other infections (e.g. chloramphenicol, isoniazid, rifampicin, sulfonamides, sulfadiazine, sulfamethiazole, sulfamethoxazole-trimethoprim, sulfaphenazole, sulfisoxazole, doxycycline and ciprofloxacin)
- Medicines used for stomach ulcers (e.g. omeprazole, sucralfate, the medicines known as H₂ antagonists including cimetidine, ranitidine, famotidine and some antacids)
- Medicines used for asthma and bronchitis (e.g. theophylline)
- Medicines used for pain and inflammation (e.g. phenylbutazone, salicylates including aspirin and steroids)
- Medicines used for sleeplessness, depression and psychiatric disorders (e.g. chlordiazepoxide, clozapine, diazepam, disulfiram, fluoxetine, methylphenidate, paroxetine, phenothiazines, quetiapine, trazodone, tricyclic antidepressants, fluvoxamine, sertraline and viloxazine)
- Medicines used for diabetes (e.g. tolbutamide)
- Medicines used for cancer (e.g. antineoplastic agents e.g. teniposide, fluorouracil), capecitabine, bleomycin, carboplatin, cisplatin, doxorubicin and methotrexate)
- Some hormone replacement therapies (oestrogens), oral contraceptives (the birth control pill) (see Pregnancy, contraception in women, and breast-feeding)
- Medicines used for organ and tissue transplants, to prevent rejection (e.g. ciclosporin, tacrolimus)
- Medicines used to lower high blood cholesterol and triglycerides (e.g. atorvastatin, fluvastatin, simvastatin)
- Medicines used in the treatment of HIV infection (e.g. delavirdine, efavirenz, fosamprenavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir)
- Medicines used to expel parasitic worms from the body (e.g. albendazole, praziquantel)
- Muscle relaxants used for surgery (neuromuscular blockers), some anaesthetic drugs (e.g. halothane, methadone)
- Some products available without a prescription (e.g. folic acid, vitamin D).

Your doctor may need to test the amount of phenytoin in your blood to help decide if any of these drugs are affecting your treatment.

The herbal preparation St John's wort (*Hypericum perforatum*) should **not** be taken at the same time as this medicine. If you already take St John's wort, consult your doctor before stopping the St John's wort preparation.

If you are being fed by a tube this can affect the concentrations of phenytoin, the active ingredient of Epanutin 30 mg/5 ml Oral Suspension, in your blood. Your doctor or pharmacist will tell you how to take this medicine with your feeds.

Epanutin 30 mg/5 ml Oral Suspension may also interfere with certain laboratory tests that you may be given.

Epanutin with food, drink and alcohol

Epanutin can be taken before or after food and drink.

Drinking a lot of alcohol can also affect the concentration of phenytoin in your blood. Talk to your doctor or pharmacist for advice.

Pregnancy, contraception in women, and breast-feeding

Pregnancy

If you are pregnant, consult your doctor promptly. You should not stop taking your medicine until you have discussed this with your doctor. Stopping your medication without consulting your doctor could cause seizures which could be dangerous to you and the pregnancy. Your doctor may decide to change your treatment. Closer monitoring of your unborn child could also be considered.

Epanutin may cause birth defects. If you take Epanutin during pregnancy your baby has a higher risk of having a birth defect. Birth defects which have been reported include facial, skull, nail, finger and heart abnormalities.

If you are of childbearing age and plan to become pregnant, consult your doctor for a preconceptional visit. You should discuss your treatment options with your doctor.

If you take Epanutin during pregnancy, your baby is also at risk for bleeding problems right after birth. Your doctor may give you and your baby a medicine to prevent this. Moreover, your child should be closely monitored.

Contraception in women

If you are of childbearing age, you should discuss your treatment options and effective methods of birth control with your doctor. Epanutin may result in a failure of hormonal contraceptives, hence you should be counselled regarding the use of other effective contraceptive methods.

Breast-feeding

Epanutin passes into breast milk. You should not take Epanutin if you are breast-feeding.

Driving and using machines

Epanutin may cause dizziness or drowsiness. If you experience these symptoms, do not drive or use any tools or machinery and contact your doctor.

Epanutin contains sucrose, ethanol and the colouring agents sunset yellow (E110) and carmoisine (E122).

This medicinal product contains small amounts of ethanol (alcohol), less than 100 mg per dose.

Epanutin contains sucrose, a type of sugar. If you have been told that you have an intolerance to some sugars, contact your doctor before taking this medicinal product. This medicine may be harmful to the teeth.

This medicine contains the colouring agents sunset yellow (E110) and carmoisine (E122) that may cause allergic reactions.

3. How to take Epanutin

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Shake the bottle vigorously before you measure your dose. Always use a medicine spoon or measure.

It is best to take Epanutin at the same time each day.

Adults

The amount of Epanutin needed varies from one person to another. Most adults need between 200mg and 500mg a day (between 7 and 17 five-ml spoonfuls of Suspension) either as a single or divided dose. Occasionally higher doses are needed.

Use in Children and adolescents

Infants and children usually start on a dose that depends on their weight (5mg per day for every kg they weigh) and is given as a divided dose, twice a day. The dose is then adjusted up to a maximum of 300mg a day (10 five-ml spoonfuls of Suspension).

Elderly

Due to decreased clearance of Epanutin, lower or less frequent dosing may be needed. The dose of Epanutin for elderly patients who may be taking other medicines may also need careful consideration and adjustment by their doctor.

Kidney or liver problems

Make sure your doctor knows if you have liver or kidney problems as you may need your dose adjusted.

If you take more Epanutin than you should

Epanutin is dangerous in overdose. If you accidentally take too much Epanutin contact your doctor at once or go to the nearest hospital casualty department. Always take the labelled medicine package with you, whether there is any Epanutin left or not.

If you forget to take Epanutin

If you forget to take a dose, take it as soon as you remember unless it is time for your next dose.

Do not take a double dose to make up for a missed dose.

If you stop taking Epanutin

Do not stop taking Epanutin unless your doctor tells you to. If you suddenly stop taking this medicine you may have a seizure. Your doctor will advise you how to stop taking the medicine.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Tell your doctor **immediately** if you experience any of the following symptoms after taking this medicine. Although they are very rare, these symptoms can be serious.

- Sudden wheeziness, difficulty in breathing, swelling of eyelids, face or lips, rash or itching (especially affecting the whole body). There is a higher incidence of this in black patients.
- If you develop potentially life-threatening skin rashes that cause blistering, (this can also affect the mouth and tongue). These may be signs of a condition known as Stevens Johnson syndrome, or toxic epidermal necrolysis (TEN). These have been reported very rarely.
- If you notice bruising, fever, you are looking pale or you have a severe sore throat. These may be the first signs of an abnormality of the blood, including decreases in the number of red cells, white cells or platelets. Your doctor may take regular blood samples to test for these effects.

- Skin rash and fever with swollen glands, particularly in the first two months of treatment, as these may be signs of a hypersensitivity reaction. If these are severe and you also experience pain and inflammation of the joints this could be related to a condition called systemic lupus erythematosus.
- Skin rash, fever, swollen glands, increase in a type of white blood cell (eosinophilia) and inflammation of internal organs (liver, lungs, heart, kidneys and large intestine) as they may be signs of a hypersensitivity reaction (Drug Reaction or rash with Eosinophilia and Systemic Symptoms (DRESS)).
- If you experience confusion or have a severe mental illness, as this may be a sign that you have high amounts of phenytoin in your blood. Your doctor may test your blood to see how much phenytoin is in the blood and may change your dose.

Other side effects that may occur are:

- **Effects on your nervous system:** Unusual eye movements, unsteadiness, difficulty in controlling movements, shaking, abnormal or uncoordinated movements, slurred speech, confusion, pins and needles or numbness, drowsiness, dizziness, vertigo, sleeplessness, nervousness, twitching muscles, headaches and change in taste.
- **Effects on your skin:** skin rash including measles-like reactions which are mild.
- **Effects on your stomach and intestines:** feeling sick, being sick and constipation.
- **Effects on your blood and lymph system:** swelling of the lymph glands.
- **Effects on your liver and kidney:** inflammation of the kidneys and liver, liver damage or liver failure which can lead to death (seen as yellowing of the skin and whites of the eye).
- **Effects on your reproductive system and breasts:** changes in the shape of the penis, painful erection.
- **Effects on your hands, face and body:** changes in the hands with difficulty in straightening the fingers, changes in facial features, enlarged lips or gums, increased or abnormal body or facial hair.
- **Effects on medical tests:** increased levels of blood sugar, or decreased levels of blood calcium, phosphate, folic acid and vitamin D. If you also do not get enough vitamin D in your diet or from exposure to sunlight, you may suffer from bone pain or fractures. Taking Phenytoin may cause abnormal thyroid test results.
- **Effects on your respiratory system:** problems breathing, inflammation of the lining of the lung.
- **Effects on your immune system:** problems with the body's defence against infection, inflammation of the wall of the arteries and immunoglobulin abnormalities.
- **Effects on your bones:** there have been reports of bone disorders including osteopenia and osteoporosis (thinning of the bone) and fractures. Check with your doctor or pharmacist if you are on long-term antiepileptic medication, have a history of osteoporosis, or take steroids.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report any side effects directly (see details below). By reporting side effects you can help provide more information on the safety of this medicine.

United Kingdom

Yellow Card Scheme website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

Malta

ADR Reporting website: www.medicinesauthority.gov.mt/adrportal

5. How to store Epanutin

Keep this medicine out of the sight and reach of children.

Do not store above 25°C

Do not use this medicine after the expiry date which is printed on the bottle label after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Epanutin 30 mg/5 ml Oral Suspension contains

Each 5 ml dose contains 30 mg of the active ingredient phenytoin.

The other ingredients are aluminium magnesium silicate, sodium benzoate (E211), citric acid monohydrate, carmellose sodium, glycerol, polysorbate 40, sucrose, ethanol (each 5 ml dose contains 0.493% of ethanol), vanillin, banana flavour, orange oil, carmoisine (E122), sunset yellow (E110) and purified water.

What Epanutin looks like and contents of the pack

Epanutin 30 mg/5 ml Oral Suspension is a cherry red liquid and is available in bottles containing 500 ml of suspension.

Marketing Authorisation Holder

Pfizer Limited
Ramsgate Road
Sandwich
Kent
CT13 9NJ
United Kingdom

Manufacturer

Delpharm Orleans
5 Avenue de Concyr
45071 Orléans
Cedex 02
France

Company Contact Address

For further information on this medicine please contact Medical Information at Pfizer Limited, Walton Oaks, Tadworth, Surrey, KT20 7NS; Tel 01304 616161.

This leaflet was last revised in: 12/2019

Ref: EP 25_0