

Name of Trust / logo

Memo (for local adaptation) - shortage of supply

To:
From:
Date:

Re: Diamorphine injection 5 and 10mg

Description of product affected

Diamorphine is licensed for the treatment of severe pain associated with surgical procedures, myocardial infarction or pain in the terminally ill and for the relief of dyspnoea in acute pulmonary oedema.¹ It is also used for labour pain in obstetrics and in women having a Caesarean Section.² In addition, a small number of people in the UK may be receiving diamorphine for the management of opioid addiction.³

Background

Accord have ongoing issues at their manufacturing plant in Europe and have been unable to support the UK with stock of diamorphine 5mg and 10mg injection, since summer 2019, this is not expected to resolve until summer 2020. Wockhardt, the only other UK supplier, have partially been supporting the UK market, with secondary care having to significantly reduce usage to ensure enough stock remains for primary care. Wockhardt experienced a delivery delay of their active pharmaceutical ingredient (API) and will also go out of stock of diamorphine 5mg from early March and 10mg from mid-March 2020. The anticipated resolution date for the Wockhardt out of stock is early April 2020. Both primary and secondary care will have to prepare for a complete out of stock of diamorphine 5mg and 10mg injection. Other strengths of diamorphine injection are not affected but there is not enough stock of these to support an increase in demand.

Alternative agents and management options

The UK is the only country that uses diamorphine for medicinal analgesic purposes. Diamorphine is metabolised to morphine and in terms of analgesic efficacy and effect on mood, it has no clinical advantages over morphine by oral or subcutaneous/intramuscular routes.⁴ In addition, morphine injection is less costly than diamorphine and does not have to be reconstituted. NICE guidance on the effective prescribing of strong opioids for pain in palliative care recommends initiating subcutaneous opioids with the lowest acquisition cost.⁵ However, diamorphine is much more water soluble than morphine, and may be preferred to morphine in the very few patients where high dose injections are needed, as smaller volumes can be used.⁴

NICE however do support the preferential use of intrathecal or epidural diamorphine over morphine for women after a Caesarean section.² They also recommend that for women who have had intrathecal opioids, there should be a minimum hourly observation of respiratory rate, sedation and pain scores for at least 12 hours for diamorphine and 24 hours for morphine.

Action

Morphine 10mg injection is available and preservative-free versions are available for epidural use.

Clinicians will need to decide whether morphine or another opioid is most appropriate for each patient. Care is needed when switching from one opioid analgesic to another to ensure equipotent dosage.⁶

According to BNF 3mg of diamorphine given parenterally is approximately equivalent to 5mg of morphine given parenterally.⁷ Therefore to convert diamorphine to morphine multiply the usual dose of diamorphine by 1.7 to give an approximately equivalent dose of morphine. However it is strongly recommended that any dose conversion should be carried out after consultation with the specialist involved.

Dose of parenteral diamorphine	Approx. equiv dose of parenteral morphine
2.5mg	4mg
5mg	8 mg
10mg	17mg
15mg	25mg
20mg	35mg
30mg	50mg

Patients should be carefully monitored after any drug switch and dose titration may be required. When converting from diamorphine to other subcutaneous drugs, consideration will also need to be given to drug compatibility in the syringe driver and the total volume of infused drugs.⁶ When converting to alternatives in regional anaesthesia, consideration will need to be given to use of preservative-free opioids.⁸

As morphine is not as soluble as diamorphine and the maximum concentration available is 30mg/mL, this may be an issue for patients requiring high doses of subcutaneous morphine, particularly bolus doses for breakthrough pain where the volume given should not exceed 2mL. If volume is an issue, advice should be sought from the palliative care team.⁹

Midwives administer diamorphine under Midwives Exemptions from the Medicines Act. This exemption also allows them to administer morphine.¹⁰ If stock of diamorphine run out, protocols need to be rewritten and midwives trained on administering morphine instead of diamorphine, as well as morphine added to Midwives Exemptions list on e-prescribing systems.

NICE do recommend a different duration of monitoring for morphine (24 hours) than diamorphine (12 hours) after Caesarean delivery and local guidance may need to be reviewed. In reality it has been shown that the incidence of clinically significant respiratory depression seen with epidural morphine after caesarean delivery is very low - in one systematic review it was estimated to occur in between 1.08 and 1.63 cases per 10,000 women.¹¹

Patients in drug addiction treatment programmes may experience difficulties switching to alternatives¹² and the community drug and alcohol team should be contacted for advice.

Please refer to local guidance, the BNF or the Palliative care formulary for information on dose conversion to other opioids; and contact relevant specialist teams for advice on management of individual cases.

References

1. Wockhardt UK Ltd. Diamorphine Injection BP 10mg. SPC, date of revision of text, 12/10/2015: <https://www.medicines.org.uk/emc/product/1466/smpc>
2. NICE. Caesarean Section Clinical Guideline No. 132. Published Nov. 2011, last updated Sept 2019. Available: <https://www.nice.org.uk/guidance/cg132>
3. The Maudsley. Prescribing guidelines in Psychiatry, 13th edition (published 2018)
4. Palliative Care Formulary, 6th edition (published 2017)
5. NICE. Opioids in palliative care: safe and effective prescribing of strong opioids for pain in palliative care of adults (CG140); Clinical guideline, published: 23 May 2012: <https://www.nice.org.uk/guidance/cg140/resources/palliative-care-for-adults-strong-opioids-for-pain-relief-35109564116677>
6. Royal College of Anaesthetists, Faculty of Pain Medicine. Opioids Aware: <https://www.rcoa.ac.uk/faculty-of-pain-medicine/opioids-aware>
7. BNF. Prescribing in palliative care, accessed online, 15 May 2019: <https://www.medicinescomplete.com/#/content/bnf/PHP107735?hspl=palliative&hspl=care>
8. Wockhardt UK Ltd. Morphine Sulfate 10mg/ml Injection BP: SPC, date of revision of text, 14/12/2018: <https://www.medicines.org.uk/emc/product/2244/smpc>
9. NHS Greater Glasgow and Clyde. Supply difficulties with diamorphine injection: guidance on alternatives for palliative patients (February 2005): http://live.nhsggc.org.uk/media/224583/nhsgg_palliative_care_guidelines_alternative_to_diamorphine.pdf
10. Nursing and Midwives Council. PRACTISING AS A MIDWIFE IN THE UK: An overview of midwifery regulation, published 28 March 2017, updated: 29 Jan 2019: <https://www.nmc.org.uk/globalassets/sitedocuments/nmc-publications/practising-as-a-midwife-in-the-uk.pdf>
11. Sharawi N et al. A systematic review evaluating neuraxial morphine and diamorphine-associated respiratory depression after Cesarean delivery. *Anesth Analg*. 2018; 127: 1385-95
12. Clark C. Diamorphine: the return of an old friend. *Hospital Pharmacy Europe* 2009; issue 42: <http://www.hospitalpharmacyeurope.com/christine-clark/diamorphine-return-old-friend>

Acknowledgements

- Jane Bass, Senior Pharmacist, Women's Services, Guy's and St Thomas' NHS Foundation Trust
- Steven Wanklyn, Consultant pharmacist for palliative and end of life care, Guy's and St Thomas' NHS Foundation Trust
- Christopher Meddings, Highly Specialist Pharmacist – Critical Care, Guy's and St Thomas' NHS Foundation Trust

Original document prepared by:

David Erskine and Yuet Wan. London and South East Regional Medicines Information,
10th May 2018, updated 16th May 2019, 11th November 2019 and 24th February 2020:
contact medicinesinformation@gstt.nhs.uk

Document modified by:

Name of individual at other centre using the product with modifications, centre, date

For all correspondence please contact:

Name of person at **base** hospital where memo is circulated (i.e. NOT the original author at Guy's and St Thomas' NHS Foundation Trust)

Disclaimer: This memo can be adapted for local use. The content does not reflect national guidance. Some of this memo is based on **clinical opinion** from practitioners. Users should bear this in mind in deciding whether to base their policy on this document. Individual trusts should ensure that procedures for unlicensed medicines are followed where a foreign import drug is required in the interim. Any decision to prescribe off-label must take into account the relevant GMC guidance and NHS Trust governance procedures for unlicensed medicines. Prescribers are advised to pay particular attention to the risks associated with using unlicensed medicines or using a licensed medicine off-label.